

# SAVE ON SPRING DEWORMING

February 1 through March 31, 2024

## DEALER & DVM DISCOUNT DAYS

In 2024, get expanded selection and savings on parasiticides from Norbrook®.

Including Noromectin® (ivermectin) Injection; Noromectin® Plus (ivermectin/clorsulon) Injection, the one that controls adult liver flukes; Eprizero® (eprinomectin) Pour-On, the one with zero meat withdrawal and zero milk discard; and now, Tauramox™ (moxidectin) Injectable Solution.

Mix & match any of the four listed products, in any size.

Receive up to a 6% discount on a single invoice purchase of \$2,500.

Receive up to an 8% discount on a single invoice purchase of \$7,500, which must include \$1,000 of Tauramox™.



Noromectin® (ivermectin) Injection • Noromectin® Plus (ivermectin/clorsulon) Injection  
Eprizero® (eprinomectin) Pour-On • Tauramox™ (moxidectin) Injectable Solution

ARMOR ID #	MANUF. PART #	UPC	DESCRIPTION	CASE QTY	ORDER QTY
21564	03148	5023534031482	EPRIZERO POUR ON 2.5 L	4	
21566	03218	5023534032182	EPRIZERO POUR ON 20 L	1	
21565	03149	5023534031499	EPRIZERO POUR ON 5 L	2	
23213	11080	5023534008446	NOROMECTIN 1% INJ 1000 ML	6	
23215	11078	5023534005322	NOROMECTIN 1% INJ 250 ML	12	
23214	11076	5023534005308	NOROMECTIN 1% INJ 50 ML	12	
32018	11079	5023534005339	NOROMECTIN 1% INJ 500 ML	1	
23217	11072	5023534005285	NOROMECTIN PLUS INJ 250 ML	12	
23216	11070	5023534005261	NOROMECTIN PLUS INJ 50 ML	12	
11511			NOROMECTIN PLUS INJ 500 ML		
11512			NOROMECTIN PLUS INJ 1000 ML		
			TAURAMOX 500 ML		

### IMPORTANT SAFETY INFORMATION:

Eprizero® (eprinomectin): Observe label directions. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. Do not use in calves to be processed for veal. Not recommended for use in species other than cattle. See product labeling for full product information.

Noromectin® Plus (ivermectin/clorsulon) and Noromectin® (ivermectin) Injection: Observe label directions and withdrawal times. Consult your veterinarian for assistance in the diagnosis, treatment and control of parasitism. Do not use in female dairy cattle of breeding age or in calves to be processed for veal. Do not use in unapproved species as severe reactions, including fatalities in dogs, may result. See product labeling for full product information.

Tauramox™ (moxidectin): Cattle must not be slaughtered for human consumption within 21 days of treatment. This drug is not approved for use in female dairy cattle 20 months of age or older, including dry dairy cows. Use in these cattle may cause drug residues in milk and/or in calves born to these cows. A withdrawal period has not been established for preruminating calves. Do not use in calves to be processed for veal.

Invoice discounts will be calculated and paid for reimbursements from AGDATA reported sales.

© 2024 Norbrook Laboratories Limited. The Norbrook logos, Noromectin and Eprizero are registered trademarks and Tauramox is a trademark of Norbrook Laboratories Limited.

  
**Norbrook®**  
NORBROOK.COM

004-24-029

### CONTACT US

#### Wholesale

828 19th Avenue NE  
St. Joseph, MN 56374  
800.397.6972 Fax: 320.363.4798  
wholesaleorders@armorah.com

1240 Green Valley Road  
Beaver Dam, WI 53916  
844.840.7770 Fax: 920.885.2812  
salesorders@armorah.com

