

INTRAMAMMARY PRODUCT COMPARISON


Dry Cow Treatment Options					Indications			
Product/ Manf.	Rx Required	Min. Dry Period Length	Milk	Meat	Staph aureus	Strep agalactiae	Strep dysgalactiae	Strep uberis
Albadry Plus™ (novobiocin penicillin) Zoetis		30 Days	72 Hrs	30 Days	X	X		
Dry-Clox® (cloxacillin benzathine) BIVI	X	30 Days	0 Hrs	30 Days	X	X		
go-dry® (penicillin) Hanford's US Vet		14 Days	72 Hrs	14 Days		X		
Orbenin-DC® (benzathine cloxacillin) Merck	X	28 Days	0 Hrs	28 Days	X	X		
Quartermaster (penicillin-dihydrostreptomycin) WG Critical Care	X	42 Days	96 Hrs	60 Days	X			
Spectramast® DC (ceftiofur hydrochloride) Zoetis	X	30 Days	0 Hrs	16 Days	X		X	X
ToMORROW® (cephapirin benzathine) BIVI		30 Days	72 Hrs	42 Days	X	X		

Lactating Treatment Options				Indications						Dosage
Product	Rx Required	Milk	Meat	Coagulase-neg. staph	Staph aureus	Strep agalactiae	Strep dysgalactiae	Strep uberis	E. coli	
Amoxi-Mast® (amoxicillin) Merck	X	60 Hrs	12 Days		X	X				1 syringe; may repeat every 12 hours for a maximum of 3 treatments
Dariclox® (sodium cloxacillin) Merck	X	48 Hrs	10 Days		X	X				1 syringe; may repeat every 12 hours for a maximum of 3 treatments
Pirsue® (pirlimycin hydrochloride) Zoetis	X	36 Hrs	9-21* Days		X	X	X	X		1 syringe; may repeat every 24 hours for up to 8 consecutive days
PolyMast™ (hetacillin potassium) BIVI	X	72 Hrs	10 Days		X	X	X		X	1 syringe; may repeat every 24 hours for a maximum of 3 treatments
Spectramast® LC (ceftiofur hydrochloride) Zoetis	X	72 Hrs	2 Days	X			X		X	1 syringe; may repeat every 24 hours for up to 8 consecutive days
ToDAY® (cephapirin sodium) BIVI		96 Hrs	4 Days		X	X				1 syringe every 12 hours for a maximum of 2 treatments

*9-day slaughter withhold required after last treatment when used for 2 days;
21-day slaughter withhold when treatment period has been 3-8 days.

DETECT. DEFEND. DELIVER.